



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

SOUTH THE PASSESS OF THE PASSESS OF

THE PARTY OF THE P

EXPERIENCE ANTICOLOGICAL SECURIOR SECUR



OFFICE OF NAVAL RESEARCH

Contract N00014-85-K-0558

Task No. NR413C012-01

TECHNICAL REPORT NO. 3

α-Methylstyrl- and Styrlphosphazene Monomers and Polymers

by

Christopher W. Allen and Jonathan C. Shaw

Accepted for Publication

in

Phosphorus and Sulfur

University of Vermont Department of Chemistry Burlington, VT 05405

October 3, 1986



Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

	,	4.27			
ŧ(URITY CLASSIFICATION OF THIS PAGE	As	A173702		
	The second of this rade		MENTATION PAGE		
i a	REPORT SECURITY CLASSIFICATION	TEV ON T DOCO	16. RESTRICTIVE MARKINGS		
	Unclassified		TO. RESTRICTIVE WARKINGS		
à	DECLASSIFICATION AUTHORITY DECLASSIFICATION / DOWNGRADING SCHEDULE PERFORMING ORGANIZATION REPORT NUMBER(S)		3. DISTRIBUTION/AVAILABILITY OF REPORT		
b			Available for public release. Distribution unlimited.		
_			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
4)					
	Technical Report No. 3				
a	NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZA	ATION	
	University of Vermont	(ii appiicable)	ONR		
٠,	ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (City, State, and ZIP Code)		
	Department of Chemistry		Office of Naval Researc	-	
	Cook Physical Science Buildingurlington, VT 05405	ng	Arlington, VA 22217		
a	MAME OF FUNDING/SPONSORING	8b. OFFICE SYMBOL	9. PROCUREMENT INSTRUMENT IDENT	IFICATION NUMBER	
	ORGANIZATION	(If applicable)			
١,	ONR ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS		
	Office of Naval Research		PROGRAM PROJECT TA	ASK WORK 'JUIT	
	800 N. Quincy			O NR ACCESSION NO	
1	Arlington, VA 22217 IIILE (Include Security Classification)		N00014-85 K-0558	4130012	
	PERSONAL AUTHOR(S) Christopher V TYPE OF REPORT 13b. TIME (W. Allen and Jon	athan C. Shaw 14. DATE OF REPORT (Year, Month, Day,) 15. PAGE COUNT	
	Preprint FROM_	то	1986/10/3	4	
	SUPPLEMENTARY NOTATION	ni 1 1.a			
16		Phaenhamie and C	HI fire		
16	Accepted for publication in I	thosphorus and 3	arrar		
'6 '~	COSATI CODES	18. SUBJECT TERMS	(Continue on reverse if necessary and ide	entify by block number)	
16 			(Continue on reverse if necessary and ideazenes homopolymers		
	COSATI CODES GROUP SUB-GROUP	18. SUBJECT TERMS cyclophosph arylphospha copolymers	(Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios		
.~	COSATI CODES GROUP SUB-GROUP ARCHEOL (CONTINUE on reverse if necessary)	18. SUBJECT TERMS cyclophospha arylphospha copolymers y and identify by block	(Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr		
	COSATI CODES GROUP SUB-GROUP ARCHAEL (COntinue on reverse if necessary FORDACT: The synthesis of	18. SUBJECT TERMS cyclophospha arylphospha copolymers y and identify by block the m and p x-m	(Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr	PART \ Lebilheresh = 1	
	GROUP SUB-GROUP AND THE COntinue on reverse if necessary FORDACT: The synthesis of to 1,2) is reported. At the	18. SUBJECT TERMS cyclophospha arylphospha copolymers y and identify by block the m and p to me bis stage of si	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na	P3F) \ [Chilha(ch) ch	
	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
.~	GROUP SUB-GROUP GROUP SUB-GROUP Address (Continue on reverse if necessary FORDACT: The synthesis of to 1,2) is reported. At the cis non-geminal species prec	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
.~	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
.~	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
.~	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
.~	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers and identify by block the m and p to me bis stage of selominating. The	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chi] c(ch) = 0 e observed with the ives enter into te mechanisms are	
	GROUP SUB-GROUP GROUP	18. SUBJECT TERMS cyclophospha arylphospha copolymers y and identify by block the m and p to me bis stage of sideminating. The ctions wherein bis homopolymeriza	Continue on reverse if necessary and ideazenes homopolymers zenes reactivity ratios nmr number) Pro ethylstypl-phosphazenes, Na ubstitution, all isomers ar mono d-methylstyrl derivat oth terminal and pennultima	P3F \ [Chille(ch) = 0 e observed with the ives enter into te mechanisms are lotriphosphazene	



Accession For	
NTIS COURT DitC TAX Unconserved tention	A C

Q-METHYLSTYRL-AND STYRLPHOSPHAZENE MONOMERS AND POLYMERS

" of ion/

Mahility Codes

CHRISTOPHER W. ALLEN* and JONATHAN C. SHAW Department of Chemistry, University of Vermont, Burlington, Vermont 05405, U.S.A.

Special

Abstract The synthesis of the m and p α -methylstyrl-phosphazenes, N₃P₃F_{6-n}[C₆H₄C(CH₃)=CH₂]_n (n=1,2) is reported. At the bis stage of substitution, all isomers are observed with the cis non-geminal species predominating. The mono α -methylstyrl derivatives enter into facile copolymerization reactions wherein both terminal and pennultimate mechanisms are observed. The synthesis and homopolymerization of styrlpentafluorocyclotriphosphazene is also reported.

Recent work has shown that the range of synthesis of new and interesting cyclo- and polyphosphazene derivatives can be dramatically enhanced by incorporation of an organofunctional unit as a substituent on a phosphorus atom. 1 This substituent can serve as the site for various synthetic transformations leading to new monomeric and polymeric phosphazenes. The use of the paraaminophenoxy substituent as an organofunctional unit has been explored in detail by the Allcock group² and by Kumar et. al.³ We have focused our investigations on alkeny14,5 and alkyn16 substituents. The alkenyl phosphazenes represent novel olefinic monomers which allow for the incorporation, via copolymerization, of the cyclophosphazenes and many of their useful properties into traditional organic polymers. 7 We have been conducting a systematic study of the synthesis. spectroscopic properties and reactivity of olefins directly bonded to the cyclophosphazene moiety 5,8 and those separated from the phosphorus atom by a spacer group.4,9 This particular series of investigations involves olefinic groups separated from the cyclophosphazene by a phenyl group.

The first system to be investigated was the α -methylstyrene system. Both the meta (1) and para (2) α -methylstyrylpenta-fluorocyclotriphosphazenes were prepared from hexafluorocyclotriphosphazene and the appropriate lithium reagent. All new materials were characterized by elemental analysis, mass spectrometry, ir and nmr (1 H, 13 C, 19 F, 31 P) spectroscopy. The chemical

$$N_3P_3F_6$$
 + $Li \bigcirc C(CH_3)=CH_2 \longrightarrow N_3P_3F_5 \bigcirc C(CH_3)=CH_2$

shifts of the vinylidene, =CH2 protons are identical in 1 and 2 suggesting the absence of significant mesomeric electron withdrawal from the aryl unit by the phosphazene. The difference in the B-vinyl carbon chemical shifts fall between those observed in systems with significant mesomeric interactions and those where such interactions cannot occur. This observation indicates either low degrees of mesomeric interaction or the more likely effect of electrostatic stabilization of aryl change density by the positive phosphorus center. At the stage of bis substitution, all three isomers of N₃P₃F₄|C₆H₄C(CH₃)=CH₂J₂ are formed. These systems were characterized by GC-mass spectrometry and nmr spectroscopy. The ratio of geminal to cis-non-geminal to trans-nongeminal isomers is 1:6.1:1.5 for the para-α-methylstyrl derivatives and 1:2.8:1.1 in the meta system. Although the predominance of non-geminal isomers can be ascribed to steric effects, the preferential formation of the cis isomer is counterintuitive. This observation may be rationalized by postulating an electrostatic attraction between the electron poor aryl ring on the phosphazene and the electron rich aryl ring in the lithium reagent thus favoring a cis attack. The decreased amount of the cis isomer in the meta derivative can be ascribed to the steric effect of the meta substituent.

Both 1 and 2 enter into facile copolymerization reactions with styrene or para-chlorostyrene. The phosphazene incorporation in the copolymer is the highest observed for alkenyl-

PARTICL STRUCTURE BRITAINS AND THEOREM STRUCTURE BESCHOOL STREET CONTROL STRUCTURE

phosphazene monomers. The maximum degrees of incorporation achieved requires that there are regions of at least two phosphazene units linked together. This structural feature has not been previously observed in organophosphazene copolymers. Both of these observations are related to the moderation of the phosphazene induced olefin polarity by the aryl spacer group interposed between the two functionalities in question. Semiquantitative information on the magnitude of this effect has been obtained from the reactivity ratios and Alfrey-Price (Q,e) parameters for 1 and 2. The reactivity ratios (r_2) of 1 and 2 as calculated by the Mortimer-Tidwell non-linear least squares approach are within the 95% joint confidence limits of each other suggesting similar

reactivity. The reactivity ratio for styrene (r₁) as well as r₂ are both less than one indicating an alternative tendency. This suggests the possibility of an electrostatic interaction between the nucleophilic styryl group and the electrophilic phosphazene. The calculated Q values for 1 and 2 are reduced from styrene showing a reduction in mesomeric interactions. The calculated e values are significantly higher than that of styrene or nitrostyrene reflecting the highly polar nature of 1 and 2. The thermal decomposition of the copolymers of 1 and 2 when studied by TGA show a two step unzipping process corresponding to phosphazene and styrene loss respectively as the temperature increases. The decomposition of the copolymers of 1 begins before similar event for copolymers of 2 indicating a role for steric destabilization in the meta substituted derivatives. The copolymers of 1 and 2 with methyl methacrylate were also obtained. Attempts to calculate reactivity ratios lead to solutions with very large 95% joint confidence limits in the phosphazene monomer. We were, however, able to achieve a good fit of the data using a pennultimate model. Thus, in these systems, the reactivity of the terminal radical is significantly modified by the presence of a phosphazene function in the pennultimate position.

Although a broad spectrum of copolymers could be prepared from 1 or 2, all attempts (radical, cationic, anionic) at homopolymerization were unsuccessful. We therefore turned our attention to the preparation of a phosphazene bearing a styrene, rather than an α -methylstyrene, substituent. Since anionic polymerization of styrene is initiated by organolithium reagents, indirect routes were explored. Aryl pentafluorocyclotriphosphazenes with carbonyl groups in the para position (3,4) were prepared from para-bromobenzaldehyde and para-bromoacetophenone by first converting the carbonyl to the acetal and then preparing the lithium

$$N_3P_3F_5$$
 CH (3) $N_3P_3F_5$ CCH₃ (4)

accepted respected accepted recepted because in the second

reagent. After coupling with N₃P₃F₆ the protecting group was removed to give 3 and 4. However, the Wittig reaction on 3 gave only degradation products as did the reduction of 4 to the alcohol. An alternative route was developed starting from an α -methoxystyryl phosphazene (5). Interestingly, 5 was unreactive with respect to borohydride or aluminum hydride addition. Fortunately, hydrostanation of 5 proceeded smoothly and β -elimination of tributyltinmethoxide from the resulting organotin derivative occured spontaneously on silica gel giving rise to the styrylpentafluorocyclotriphosphazene (6).

N₃P₃F₅
$$\longrightarrow$$
 C(OMe)=CH₂ \longrightarrow {N₃P₃F₅ \bigcirc CH(OMe)CH₂Sn(C₄H₉)₃}
(5) (C₄H₉)₃SnH \downarrow Silica
N₃P₃F₅ \bigcirc CH=CH₂ (6)

The ¹³C chemical shift of the vinylidene carbon in 6 occurs further downfield than even 4-nitrostyrene, making 6 amongst the most electron deficient styrene derivatives known. Polymerization of 6 via radical initiation proceeded smoothly to give the homopolymer (7), a polymer having a carbon chain backbone and an inorganic surface. The molecular weights of 7 are 102,000 and

374,000 for $M_{\rm H}$ and $M_{\rm W}$ respectively. Thermal decomposition of 7, as monitored by TGA, starts about 80° higher than that for polystyrene and the polymer retains fifteen to thirty percent of its original weight at temperatures of 900° to 1100°C. This observation suggests the possibility of thermal decomposition of carbon chain polymers with inorganic surfaces as a potentially new route to ceramic materials.

Acknowledgements

This work was supported in part by the Office of Naval Research.

References

- 1) Allen, C.W. <u>J. Polym. Sci.</u>, <u>Polym. Symp.</u> 1983, <u>70</u>, 99.
- 2) Allcock, H.R.; Hymer, W.C.; Austin, P.E. Macromolecules 1983, 16, 1401.
- 3) Kumar, D.; Fohlen, G.M.; Parker, J.A. <u>J. Polym. Sci.</u>, Polym. Chem. Ed. 1984, 22, 1411.
- 4) Ramachandran, K.; Allen, C.W. Inorg. Chem. 1983, 22, 1445.
- 5) Allen, C.W., Bright, R.P. <u>Inorg. Chem.</u> 1983, 22, 1291.
- 6) Allen, C.W., Desorcie, J.L., Ramachandran, K. J. Chem. Soc., Dalton Trans. 1984, 2843.
- 7) Dupont, J.G.; Allen, C.W. Macromolecules 1979, 12, 169.
- B) Dupont, J.G., Allen, C.W. Inorg. Chem. 1978, 17, 3093.
- 9) Allen, C.W., Bright, R.P. Inorg. Chim. Acta 1985, 99, 107.

DL/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

<u>c</u>	No. Copies		to. <u>Opics</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Mayal Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1 2	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
PINSRDC Attn: Dr. H. Singerman Applied Chemistry Division Apparatis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
On. William Tolles Superintendent Chemistry Division, Code 6100 Chaval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	l

ABSTRACTS DISTRIBUTION LIST, 356B

Contract of the Contract of th

Professor A. G. MacDiarmid
Department of Chemistry
University of Pennsylvania
Philadelphia, Pensylvania 19174

Naval Ship Research and Development Center Annapolis, Maryland 21402

Professor H. Allcock
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Professor R. Lenz
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts
01002

Professor A. Epstein
Department of Physics
Obio State University
Columbus, Obio 43212

Br. J. Griffith Continuous Research Laboratory on the continuous Chemistry Section, Code 6120 Washington, D.C. 20375-5000

Professor G. Wnek
Department of Materials Science in the department of Materials Science and Engineering
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. Samson Jenekhe Honeywell Corporate Technology Center 19701 Lyndale Avenue South Bloomington, Minnesota 55420

Professor William R. Krigbaum Department of Chemistry Duke University Ducham, North Carolina 27706

Tyan Caplan binsroc Code 0125 Annapolis, MD 21401 Dr. Richard M. Laine SRI International 333 Ravenswood Avenue Menlo Park, California 94025

Dr. L. Buckley Naval Air Development Center Code 6063 Warminster, Pennsylvania 18974

Dr. James McGrath
Department of Chemistry
Virginia Polytechnic Institute
Blacksburg, Virginia 24061

Dr. Geoffrey Lindsay Chemistry Division Naval Weapons Center China Lake, California 93555

Professor C. Allen
Department of Chemistry
University of Vermont
Burlington, Vermont 05401

Or. William Tolles Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000

Professor T. Katz Department of Chemistry Columbia University New York, New York 10027

Professor J. Salamone Department of Chemistry University of Lowell Lowell, Massachusetts 01854

Dr. B. Gordon III
Department of Materials Science
Pennsylvania State University
University Park, PA 16802

Dr. R. Miller Almaden Research Center 650 Harry Road San Jose. CA 95120-6099

ABSTRACTS DISTRIBUTION LIST, 356B

Professor T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Kurt Baum Fluorochem, Inc. 680 S. Ayon Avenue Azuza, California 91702

Professor Stephen Wellinghoff Division of Chemistry and Chemical Engineering Southwest Research Institute 6220 Culebra Road Postal Drawer 283510 San Antonio, Texas 78284

hr. William Bailey
Department of Chemistry
University of Maryland
Folloge Park, Maryland 20742

Fig. J.C.H. Chien
Department of Polymer Science and
Engineering
University of Massachusetts
Amherst, MA 01003

Professor G. Whitesides
Pepartment of Chemistry
Harvard University
Cambridge, Massachusetts 02138

Ur. K. Paciorek Ultrasystems, Inc. P.O. Box 19605 Irvine, California 92715

がある。 1990年の大学のでは、これのでは、1990年のできたが、1990年のでものできたが、1990年のでものでものでものできたが、1990年のでものできたが、1990年のできたが、1990年のできたが、1990年のできたが、199

Br. Ronald Archer
Pepartment of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

Professor D. Seyferth
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Professor J. Moore Department of Chemistry Rensselaer Polytechnic Institute Troy, New York 12181

Dr. V. Percec Department of Macromolecular Science Case Western Reserve University Cleveland, Ohio 44106

Dr. Gregory Girolami Department of Chemistry University of Illinois Urbana-Champagne, IL 61801

Dr. Ted Walton Chemistry Division Code 6120 Naval Research Lab Washinston D.C. 20375-5000

Dr. Thomas J. McCarthy
Polymer Science and
Engineering Department
University of Massachusetts
Amherst, MA 01003

Professor A. Heeger Department of Physics University of California Santa Barbara, California 93106

Dr. Fred Wudl
Department of Chemistry
University of California
Santa Barbara, CA 93106